

Name : _____

Grade : VIII

Subject : Mathematics

Chapter : 12. Exponents and Powers

Objective Type Questions

1 Marks.

I. Multiple choice questions

1. The multiplicative inverse of $\left(\frac{-5}{9}\right)^{-99}$ is [NCERT Exemplar]
 a. $\left(\frac{-5}{9}\right)^{-99}$ b. $\left(\frac{5}{9}\right)^{-99}$ c. $\left(\frac{9}{-5}\right)^{-99}$ d. $\left(\frac{9}{5}\right)^{99}$
2. What is the value of 'm' if $(-2)^2 \times (-5)^2 = 50m$?
 a. 10 b. 2 c. 100 d. -100
3. What is the scientific notation of 0.0023?
 a. 2.3×10^{-3} b. 23×10^{-3} c. 2.3×10^3 d. 23×10^3
4. The usual form for 2.03×10^{-5} : [NCERT Exemplar]
 a. 0.203 b. 0.00203 c. 203000 d. 0.0000203
5. $\left(\frac{1}{10}\right)^0$ is equal to:
 a. 0 b. $\frac{1}{10}$ c. 1 d. 10
6. $\left(\frac{-7}{5}\right)^{-1}$ is equal to: [NCERT Exemplar]
 a. $\frac{5}{7}$ b. $\frac{-5}{7}$ c. $\frac{7}{5}$ d. $\frac{-7}{5}$
7. The value of $(7^{-1} - 8^{-1})^{-1} - (3^{-1} - 48^{-1})^{-1}$ is:
 a. 44 b. 56 c. 68 d. 12
8. $(2^{-1} + 3^{-1} + 4^{-1})^{-1}$ is:
 a. $\frac{1}{2} + \frac{1}{3} + \frac{1}{4}$ b. $2 + 3 + 4$ c. $1 + 1 + 1$ d. 1
9. $\left(\frac{3}{4}\right)^5 \div \left(\frac{5}{3}\right)^5$ is equal to:
 a. $\left(\frac{3 \div 5}{5 \div 3}\right)^5$ b. $\left(\frac{3}{4} \div \frac{5}{3}\right)^1$ c. $\left(\frac{3}{4} \div \frac{5}{3}\right)^0$ d. $\left(\frac{3}{4} \div \frac{5}{3}\right)^{10}$
10. $(-7^3 \div 7^{-8}) \div 7^5 =$
 a. 7^6 b. 7^{-6} c. -1 d. 1

11. The value of p , for which $7^7 \div 7^{-p} = 7^{10}$, is :
 a. 1 b. 2 c. 3 d. 4
12. When $x = 2$, $x(x^x) - x$ is equal to
 a. 4 b. 6 c. 8 d. 10
13. Let x be any non-zero integer and m, n be negative integer. The $x^m \times x^n$ is equal to
 [NCERT Exemplar]
 a. x^m b. x^{m+n} c. x^n d. x^{m-n}
14. The standard form for 0.000064 is:
 [NCERT Exemplar]
 a. 64×10^4 b. 64×10^{-4} c. 6.4×10^5 d. 6.4×10^{-5}
15. $(-9)^3 \div (-9)^8$ is equal to
 [NCERT Exemplar]
 a. $(9)^5$ b. $(9)^{-5}$ c. $(-9)^5$ d. $(-9)^{-5}$
16. For a non-zero rational number z , $(z^{-2})^3$ is equal to:
 [NCERT Exemplar]
 a. z^6 b. z^{-6} c. z^1 d. z^4
17. Which of the following is not the reciprocal of $(\frac{2}{3})^4$?
 a. $(\frac{3}{2})^4$ b. $(\frac{3}{2})^{-4}$ c. $(\frac{2}{3})^4$ d. $\frac{3^4}{2^4}$
18. In 2^2 , n is known as
 [NCERT Exemplar]
 a. base b. constant c. exponent d. variable
19. Cube of $-\frac{1}{2}$ is:
 a. $-\frac{1}{8}$ b. $\frac{1}{16}$ c. $-\frac{1}{8}$ d. $-\frac{1}{16}$
20. For a non-zero integer x , $x^7 \div x^{12}$ is equal to:
 a. x^5 b. x^{19} c. x^{-5} d. x^{-19}

1. c	2. b	3. a	4. d	5. c	6. b	7. a	8. d	9. a	10. c
11. c	12. b	13. b	14. d	15. d	16. b	17. b	18. c	19. c	20. c

II. Multiple choice questions

1. Multiplicative inverse of 2^7 is
 [NCERT Exemplar]
 a. 2^{-7} b. 7^2 c. -2^7 d. 2^7
2. Then value of $\frac{1}{4^{-2}}$ is
 [NCERT Exemplar]
 a. 16 b. 8 c. $\frac{1}{16}$ d. $\frac{1}{8}$

3. Then reciprocal of $\left(\frac{2}{5}\right)^{-1}$ is

[NCERT Exemplar]

- a. 16 b. 8 c. $\frac{1}{16}$ d. $\frac{1}{8}$

4. If y be any non-zero integer, then y^0 is equal to

[NCERT Exemplar]

- a. 1 b. 0 c. -1 d. Not defined

5. Which of the following is equal to $\left(-\frac{3}{4}\right)^{-3}$?

- a. $\left(\frac{3}{4}\right)^{-3}$ b. $\left(-\frac{4}{3}\right)^{-3}$ c. $\left(\frac{4}{3}\right)^3$ d. $\left(-\frac{4}{3}\right)^3$

6. The value of $(7^{-1} - 8^{-1})^{-1} - (3^{-1} - 4^{-1})^{-1}$ is

[NCERT Exemplar]

- a. 44 b. 56 c. 68 d. 12

7. The standard form for 0.000064 is

[NCERT Exemplar]

- a. 64×10^4 b. 6.4×10^5
c. 6.4×10^{-5} d. None of these

8. The usual form for 2.03×10^{-5}

- a. 0.203 b. 0.00203 c. 203000 d. 0.0000203

9. $\left[\left\{\left(-\frac{1}{2}\right)^2\right\}^{-2}\right]^{-1}$ is equal to

- a. 16 b. -16 c. $-\frac{1}{16}$ d. $\left(\frac{3}{7}\right)^{-6}$

10. If $x = \left(\frac{3}{7}\right)^{-3}$, then x^{-2} equals

- a. $\left(\frac{3}{7}\right)^{-3}$ b. $\left(\frac{3}{7}\right)^9$ c. $\left(\frac{3}{7}\right)^6$ d. $\left(\frac{3}{7}\right)^{-6}$

1. a	2. a	3. a	4. a	5. d	6. a	7. c	8. d	9. d	10. c
------	------	------	------	------	------	------	------	------	-------

I. Fill in the blanks

1. The standard form of 12340000 is _____.

[NCERT Exemplar]

2. The usual form of 3.41×10^6 is _____.

[NCERT Exemplar]

3. If $36 = 6 \times 6 = 6^2$, then $\frac{1}{36}$ expressed as a power with the base 6 is _____.

[NCERT Exemplar]

4. The value of $\left(\frac{1}{2^3}\right)^3$ is equal to _____.

[NCERT Exemplar]

5. By multiplying $(10)^5$ by $(10)^{-10}$, we get _____.

[NCERT Exemplar]

1. 1.234×10^7	2. 3410000	3. 6^{-2}	4. $\frac{1}{2^6}$	5. 6^{-2}
------------------------	------------	-------------	--------------------	-------------

I. True or False

1. The multiplicative inverse of $\left(\frac{3}{2}\right)^2$ is not equal to $\left(\frac{2}{3}\right)^{-2}$ [NCERT Exemplar]
2. $(-5)^{-2} \times (-9)^{-3} = (-5)^{-6}$. [NCERT Exemplar]
3. The multiplicative inverse of $(-4)^{-2}$ is $(4)^{-2}$. [NCERT Exemplar]
4. $\frac{x^m}{y^m} = \left(\frac{y}{x}\right)^{-m}$ [NCERT Exemplar]
5. $(-4)^{-4} \times (-7)^{-4} = (-7)^{-2}$ [NCERT Exemplar]

1. True	2. False	3. False	4. True	5. False
---------	----------	----------	---------	----------

I. Very short answer type questions.

1. Write 0.000005678 in the standard form. [NCERT Exemplar]

Sol. $0.000005678 = \frac{5678}{1000000000} = \frac{5.678}{10^9} \times 10^3$
 $= 5.678 \times 10^{-6}$

2. What is the usual form for 2.3×10^{-10} ? [NCERT Exemplar]

Sol. $2.3 \times 10^{-10} = 23 \times 10^{-1} \times 10^{-10}$
 $= 23 \times 10^{-11}$
 $= \frac{23}{100000000000}$
 $= 0.0000000023$

3. Express 16^{-2} as a power with the base 2. [NCERT Exemplar]

Sol. $16^{-2} = \frac{1}{(16)^2} = \frac{1}{(2^4)^2} = \frac{1}{2^8}$
 $[\because (a^m)^n = a^{mn}]$
 $= 2^{-8}$

4. Write 39,00,00,000 in the standard form. [NCERT Exemplar]

Sol. $39,00,00,000 = 3.9 \times 10^8$

5. Divide 293 by 10,00,000 and express the result in standard form. [NCERT Exemplar]

Sol. $\frac{293}{1000000} = \frac{2.93 \times 10^2}{10^6}$

6. Express each of the following in standard form: [NCERT Exemplar]

a. The mass of a proton in gram is

1673

b. A Helium atom has a diameter of 0.000000022 cm.

Sol. a.

$$\frac{1673}{1000000000000000000000000000}$$
$$= \frac{1.673 \times 10^3}{10^{27}}$$
$$= 1.673 \times 10^{-24} \text{ gm}$$

b. $0.000000022 = 2.2 \times 10^{-8} \text{ cm}$

7. Simplify and write in exponential form:

a. $(-2)^{-3} \times (-2)^{-4}$

b. $p^3 \times p^{-10}$

c. $3^2 \times 3^{-5} \times 3^6$

Sol. a. $(-2)^{-3} \times (-2)^{-4} = (-2)^{(-3)+(-4)}$

$$[a^m \times a^n = a^{m+n}]$$

$$= (-2)^{-7} \text{ or } \frac{1}{(-2)^7}$$

b. $p^3 \times p^{-10} = (p)^{3+(-10)} = (p)^{-70}$ or $\frac{1}{(p)^7}$

c. $3^2 \times 3^{-5} \times 3^6 = 3^{2+(-5)+6}$
 $3^{8-5} = 3^3$

8. Find the product of the cube of (-2) and the square of $(+4)$.

[NCERT Exemplar]

Sol. $(2)^3 \times (+4)^2 = (-8) \times (16)$
 $= -128$

9. Simplify: $(2^5 \div 2^8) \times 2^{-7}$.

Sol. $(2^5 \div 2^8) \times 2^{-7} = \left(\frac{2^5}{2^8}\right) \times 2^{-7}$
 $= (2^{-3}) \times 2^{-7} = 2^{-10}$

10. Express $3^{-5} \times 3^{-4}$ as a power of 3 with positive exponent.

[NCERT Exemplar]

Sol. $3^{-5} \times 3^{-4} = 3^{-9} = \frac{1}{3^9}$

II. Very short answer type questions.

1. Find the value of x for the expression $3^{5x-1} \div 27 = 3^{-5}$

Sol. $3^{5x-1} \div 27 = 3^{-5}$

$$\Rightarrow 3^{5x-1} \div 27 = 3^{-5}$$

$$\Rightarrow \frac{3^{5x-1}}{3^3} = 3^{-5} \quad \Rightarrow \quad 3^{5x-1-3} = 3^{-5}$$

$$\Rightarrow \frac{3^{5x-4}}{3^3} = 3^{-5}$$

Comparing exponent of 3

$$5x - 4 = -5$$

$$5x - 4 = -5 \quad \Rightarrow \quad 5x = -1 \quad \Rightarrow \quad x = \frac{-1}{5}$$

2. What is the value of k if 385600000 is written in the form $k \times 10^n$ with $n = 7$?

Sol. $x = 385600000 \Rightarrow x = 38.56 \times 10^7$

Hence, the value of $k = 38.56$.

3. What is the value of $(6^{-1} + 8^{-1} + 12^{-1})^0$?

Sol. As we know that 0 exponent of any base equals 1.

Hence, $(6^{-1} + 8^{-1} + 12^{-1})^0 = 1$

4. Express 300970000 in standard form.

Sol. Standard form of 300970000 is $= 3.0097 \times 10^8$

5. What is the reciprocal of $\left(\frac{8}{5}\right)^{-4}$?

Sol. For finding reciprocal we just reverse the fraction

$\therefore$ Its reciprocal $= \left(\frac{5}{8}\right)^{-4} = \left(\frac{8}{5}\right)^4$

6. What is the multiplicative inverse of $\left(-\frac{5}{9}\right)^{-99}$?

Sol. Let the multiplicative inverse be x .

then $\left(-\frac{5}{9}\right)^{-99} \times x = 1$ [$\because 1$ is identity]

$$x = \left(-\frac{5}{9}\right)^{99}$$

7. What is the usual form for 2.3×10^{-10} ?

Sol. Its usual form will be 0.00000000023.

8. What is the expression for 4^{-3} as a power with the base 2?

Sol. $(4)^{-3} = (2^2)^{-3}$

$= (2)^{-6}$, which is the required expression.

9. Express 16^{-2} as a power with the base 2.

[NCERT Exemplar]

Sol. $16 = \frac{1}{16^2} = \frac{1}{(2^4)^2} = \frac{1}{2^8}$

I. Short answer type questions.

1. Simply: $\left(\frac{1}{4}\right)^{-2} + \left(\frac{1}{2}\right)^{-2} + \left(\frac{1}{3}\right)^{-2}$

Sol. $\left(\frac{1}{4}\right)^{-2} + \left(\frac{1}{2}\right)^{-2} + \left(\frac{1}{3}\right)^{-2} = \frac{1}{\left(\frac{1}{4}\right)^2} + \frac{1}{\left(\frac{1}{2}\right)^2} + \frac{1}{\left(\frac{1}{3}\right)^2}$

$$= \frac{1}{\frac{1}{16}} + \frac{1}{\frac{1}{4}} + \frac{1}{\frac{1}{9}}$$

$$= 16 + 4 + 9$$

$$= 29$$

2. Simplify: $\frac{(-2)^3 \times (-2)^7}{3 \times 4^6}$

Sol. $\frac{(-2)^3 \times (-2)^7}{3 \times 4^6} = \frac{(-2)^{3+7}}{3 \times (2^2)^6}$

$$\{a^m \times a^n = a^{m+n}\}$$

$$= \frac{(-2)^{10}}{3 \times 2^{12}} \{(a^m)^n \times a^{m \times n}\}$$

$$= \frac{(-2)^{10}}{3 \times 2^{12}} = \frac{2^{10-12}}{3}$$

$$\{a^m \times a^n = a^{m-n}, (-2)^{10} = 2^{10}\}$$

$$\frac{2^{-2}}{2} = \frac{1}{3 \times 2^2} = \frac{1}{12}$$

3. Find the multiplicative inverse of $(-7)^{-2} \div (90)^{-1}$

Sol. $(-7)^{-2} \div (90)^{-1} = \frac{1}{(-7)^2} \div \frac{1}{90}$

$$= \frac{1}{49} \div \frac{1}{90}$$

$$= \frac{90}{49}$$

Multiplicative inverse of $(-7)^{-2} \div (90)^{-1}$

4. Evaluate: $(5^{-1} \times 2^{-1}) \times 6^{-1}$

Sol. $(5^{-1} \times 2^{-1}) \times 6^{-1} = (5 \times 2)^{-1} \times 6^{-1}$

$$= (10)^{-1} \times 6^{-1}$$

$$= (10 \times 6)^{-1}$$

$$= (60)^{-1} = \frac{1}{60}$$

5. Find the value of x , so that

Sol. $\left(\frac{5}{3}\right)^{-2} \times \left(\frac{5}{3}\right)^{-14} = \left(\frac{5}{3}\right)^{8x}$

$$\left(\frac{5}{3}\right)^{-2} \times \left(\frac{5}{3}\right)^{-14} = \left(\frac{5}{3}\right)^{8x}$$

$$\left(\frac{5}{3}\right)^{-2+(-14)} = \left(\frac{5}{3}\right)^{8x} \quad \therefore a^m \times a^n = a^{m+n}$$

$$\left(\frac{5}{3}\right)^{-2+(-14)} = \left(\frac{5}{3}\right)^{8x}$$

$$\left(\frac{5}{3}\right)^{-16} = \left(\frac{5}{3}\right)^{8x}$$

$$-16 = 8x$$

$$x = -2$$

6. Express $\frac{1.5 \times 10^6}{2.5 \times 10^{-4}}$ in the standard form.

[NCERT Exemplar]

Sol. $\frac{1.5 \times 10^6}{2.5 \times 10^{-4}} = 0.6 \times 10^{10}$

$$= 6 \times 10^{-1} \times 10^{10}$$

$$= 6 \times 10^9$$

7. Find the value of $n^{\frac{2^n \times 2^6}{2^{-3}}} = 2^{18}$

Sol. $\frac{2^n \times 2^6}{2^{-3}} = 2^{18}$

$$2^n \times 2^6 = 2^{18} \times 2^{-3}$$

$$2^{n+6} = 2^{18-3}$$

$$2^{n+6} = 2^{15}$$

$$n+6 = 15$$

$$n = 9$$

8. Find x , so that $(-5)^{x+1} \times (-5)^5 = (-5)^7$ [NCERT Exemplar]

Sol. $(-5)^{x+1} \times (-5)^5 = (-5)^7$

$$(-5)^{x+1+5} = (-5)^7 \quad [a^m \times a^n = a^{m+n}]$$

$$(-5)^{x+6} = (-5)^7$$

On both sides, powers have the same base, so their exponents must be equal.

Therefore, $x+6 = 7$

$$x = 7 - 6 = 1$$

$$x = 1$$

9. If $5^{2x+1} \div 25 = 125$, find the value of x .

Sol. Since, $\frac{5^{2x+1}}{25} = 125$

Then $\frac{5^{2x+1}}{5^2} = 5^3$

or $5^{2x+1} \times 5^{-2} = 5^3$

or $5^{2x+1} \times 5^{-2} = 5^3 \quad \left[\because \frac{1}{x^n} = x^{-n} \right]$

or $5^{2x-1} = 5^3 \quad \left[\because x^m \times x^n = x^{m+n} \right]$

Comparing powers

$$2x - 1 = 3$$

or $2x = 3 + 1$

or $2x = 4$

or $x = \frac{4}{2} = 2$

Hence, $x = 2$

10. By what number should $\left(\frac{1}{2}\right)^{-1}$ be multiplied to that the product is $\left(\frac{-5}{4}\right)^{-1}$?

Sol. Let, the number = x

According to problem,

$$x \times \left(\frac{1}{2}\right)^{-1} = \left(\frac{-5}{4}\right)^{-1}$$

or $x \times \frac{2}{1} = \frac{4}{-5}$

or $2x = \frac{-4}{5}$

or $x = \frac{-4}{5} \times \frac{1}{2} = \frac{-2}{5}$

Hence, required number = $-\frac{2}{5}$

II. Very short answer type questions.

1. Express $\frac{16}{81}$ and $\frac{-8}{27}$ as powers of a rational number.

[NCERT Exemplar]

Sol. $\frac{16}{81} = \frac{2^4}{3^4} = \left(-\frac{2}{3}\right)^3$

and $\frac{-8}{27} = \frac{(-2)^3}{3^3} = \left(\frac{2}{3}\right)^3$

2. Express as a power of a rational number with negative exponent $\left[\left(\frac{-3}{2}\right)^{-2}\right]^{-3}$

[NCERT Exemplar]

Sol. $\left[\left(\frac{-3}{2}\right)^{-2}\right]^{-3}$

$$= \left(\frac{-3}{2}\right)^{-2 \times -3} = \left(\frac{-3}{2}\right)^6$$

But we have to express the power with negative sign.

Hence, it will become $\left(\frac{-2}{3}\right)^{-6}$.

3. Find the value of x so that $\left(\frac{5}{3}\right)^{-2} \times \left(\frac{5}{3}\right)^{-14} = \left(\frac{5}{3}\right)^{8x}$.

[NCERT Exemplar]

Sol. $\left(\frac{5}{3}\right)^{-2} \times \left(\frac{5}{3}\right)^{-14} = \left(\frac{5}{3}\right)^{8x}$

$$\left(\frac{5}{3}\right)^{-2-14} \times \left(\frac{5}{3}\right)^{8x}$$

$$\left(\frac{5}{3}\right)^{-16} = \left(\frac{5}{3}\right)^{8x}$$

$$8x = -16$$

$$x = \frac{-16}{8} = -2.$$

$$[a^m \times a^n = a^{m+n}]$$

(Comparing the exponents)

4. Find the multiplicative inverse of $(-7)^{-2} \div (90)^{-1}$

[NCERT Exemplar]

Sol. $(-7)^{-2} \div (90)^{-1}$

$$\frac{1}{(7)^2} \div \frac{1}{90}$$

$$= \frac{1}{49} \times \frac{90}{1} = \frac{90}{49}$$

$$[a^{-m} = \frac{1}{a^m}]$$

Hence, the multiplicative inverse will be the reciprocal of it i.e. $\frac{49}{90}$.

5. Express $\frac{1.5 \times 10^6}{2.5 \times 10^{-4}}$ in the standard form.

[NCERT Exemplar]

Sol. $\frac{1.5 \times 10^6}{2.5 \times 10^{-4}} = \frac{1.5}{2.5} \times 10^6 \times 10^4$
 $= 0.6 \times 10^{10} = 6.0 \times 10^9$

6. Express each of the following in standard form.

[NCERT Exemplar]

i. The mass of a proton in gram is

1673

100000000000000000000000000000000

ii. Mass of a molecule of hydrogen gas is about

1673

100000000000000000000000000000000

Sol. i. $\frac{1673}{1000000000000000000000000}$

$= \frac{1.673 \times 1000}{10^{27}} = 1.673 \times 10^{-24} g$

[illegible]

7. If $a = -1, b = 2$, then find the value of the following.

[NCERT Exemplar]

i. $a^b - b^a$

ii. $a^b \div b^a$

Sol. i. $a^b - b^a = (-1)^2 - (2)^{-1} = 1 - \frac{1}{2} = \frac{1}{2}$

ii. $a^b \div b^a = (-1)^2 \div (2)^{-1}$
 $= 1 \div \frac{1}{2} = 1 \times \frac{2}{1} = 2$

8. By what number should we multiply $(-29)^0$ so that the product becomes $(+29)^0$?

[NCERT Exemplar]

Sol. Since any base having exponent 0 is equal to 1

$$\therefore (-29)^0 \text{ and } (+29)^0 = 1$$

$\Rightarrow$ 1 and product of a number equal 1

$$\therefore 1 \times 1 = 1$$

Hence 1 is the required number.

9. Express $\frac{400}{3969}$ in exponential form.

Sol. $400 = 2 \times 2 \times 2 \times 2 \times 5 \times 5 = 4^2 \times 5^2$
 $3696 = 3 \times 3 \times 3 \times 3 \times 7 \times 7 = 9^2 \times 7^2$

$$\begin{aligned}\therefore \frac{400}{3969} &= \frac{4^2 \times 5^2}{9^2 \times 7^2} = \frac{(4 \times 5)^2}{(9 \times 7)^2} \\ &= \left(\frac{4 \times 5}{9 \times 7}\right)^2 = \left(\frac{20}{63}\right)^2\end{aligned}$$

I. Long answer type questions.

1. Simplify:

a. $\left(\left(\frac{2}{3}\right)^3\right)^3 \times \left(\frac{1}{2}\right)^{-4} \times 3^{-1} \times \frac{1}{6}$

b. $\frac{49 \times z^{-3}}{7^3 \times 10 \times z^{-5}} \quad (z \neq 0)$

[NCERT Exemplar]

Sol. a. $\left(\left(\frac{2}{3}\right)^3\right)^3 \times \left(\frac{1}{2}\right)^{-4} \times 3^{-1} \times \frac{1}{6}$

$$\begin{aligned}&= \left(\left(-\frac{3}{3}\right)^2\right)^3 \times (3) \times \frac{1}{3} \times \frac{1}{6} \\ &= \left(\frac{9}{4}\right)^3 \times 81 \times \frac{1}{3} \times \frac{1}{6} \\ &= \frac{729}{64} \times 81 \times \frac{1}{3} \times \frac{1}{6} \\ &= \frac{243 \times 27}{128} = \frac{(3)^5 \times (3)^3}{2^7} = \frac{3^8}{2^7}\end{aligned}$$

b. $\frac{49 \times z^{-3}}{7^{-3} \times 10 \times z^{-5}} \quad (z \neq 0)$

$$\begin{aligned}&= \frac{7^2 \times 7^3 \times z^{-3} \times z^5}{10} \\ &= \frac{7^{2+3} \times z^{5-3}}{10} = \frac{7^5 \times z^2}{10}\end{aligned}$$

2. Find x , so that $\left(\frac{2}{9}\right)^3 \times \left(\frac{2}{9}\right)^{-6} = \left(\frac{2}{9}\right)^{2x-1}$

Sol. $\left(\frac{2}{9}\right)^3 \times \left(\frac{2}{9}\right)^{-6} = \left(\frac{2}{9}\right)^{2x-1}$

$$\left(\frac{2}{9}\right)^{3+(-6)} = \left(\frac{2}{9}\right)^{2x-1} \quad [a^m \times a^n = a^{m+n}]$$

$$\left(\frac{2}{9}\right)^{-3} = \left(\frac{2}{9}\right)^{2x-1}$$

$$-3 = 2x - 1$$

$$-3 + 1 = 2x$$

$$2x = -1$$

$$x = -\frac{1}{2}$$

3. If $\frac{5^m \times 5^3 5^{-2}}{5^{-5}} = 5^{12}$, find m .

Sol. $\frac{5^m \times 5^3 5^{-2}}{5^{-5}} = 5^{12}$

$$5^m \times 5^3 \times 5^{-2} \times 5^5 = 5^{12}$$

$$5^m \times 5^{3-2+5} = 5^{12}$$

$$5^m \times 5^6 = 5^{12}$$

$$m + 6 = 12 - 6$$

$$m = 6$$

4. Simplify: $\left(\frac{1}{5}\right)^{45} \times \left(\frac{1}{5}\right)^{-60} - \left(\frac{1}{5}\right)^{+28} \times \left(\frac{1}{5}\right)^{-43}$

Sol. $\left(\frac{1}{5}\right)^{45} \times \left(\frac{1}{5}\right)^{-60} - \left(\frac{1}{5}\right)^{+28} \times \left(\frac{1}{5}\right)^{-43}$
 $\Rightarrow \left(\frac{1}{5}\right)^{45-60} - \left(\frac{1}{5}\right)^{28-43}$
 $\Rightarrow \left(\frac{1}{5}\right)^{45-60} - \left(\frac{1}{5}\right)^{28-43}$
 $\Rightarrow \left(\frac{1}{5}\right)^{-15} - \left(\frac{1}{5}\right)^{-15}$
 $\Rightarrow (5)^{15} - (5)^{-15}$
 $\Rightarrow 0$

5. By what number should $(-8)^{-3}$ be multiplied so that the product may be equal to $(-6)^{-3}$?

Sol. Let the number be x

$$\begin{aligned} (-8)^{-3} \times x &= (-6)^{-3} \\ \left(\frac{1}{-8}\right)^3 \times x &= \left(\frac{1}{6}\right)^3 \\ \frac{1}{512} \times x &= \frac{1}{216} \\ x &= \frac{512}{216} \\ x &= \frac{64}{27} \end{aligned}$$

6. Simplify: $\frac{(3^{-2})^2 \times (5^2)^2 \times (t^{-3})^2}{(3^{-2})^5 \times (5^3)^{-2} \times (t^{-4})^3}$

Sol. $\frac{(3^{-2})^2 \times (5^2)^2 \times (t^{-3})^2}{(3^{-2})^5 \times (5^3)^{-2} \times (t^{-4})^3}$
 $= \frac{(3)^{-4} \times (5)^{-6} \times (t)^{-6}}{(3)^{-10} \times (5)^{-6} \times (t)^{-12}}$
 $= (3)^{-4} \times (3)^{10} \times (5)^{-6} \times (5)^6 \times (t)^{-6} \times (t)^{12}$
 $= (3)^6 \times (5)^{-6+6} \times t^{-6+12}$
 $= (3)^6 \times (5)^0 \times (t)^6$
 $= 729 t^6$

7. Simplify : $\frac{2^{-5} \times 3^{-5} \times 125}{5^{-4} \times 6^{-5}}$

Sol. Since, $125 = 5 \times 5 \times 5 = 5^3$

$$6^{-5} = (2 \times 3)^{-5} = 2^{-5} \times 3^{-5}$$

$$\therefore \frac{2^{-5} \times 3^{-5} \times 125}{5^{-4} \times 6^{-5}} = \frac{2^{-5} \times 3^{-5} \times 5^3}{5^{-4} \times 2^{-5} \times 3^{-5}}$$

$$= \frac{2^{-5}}{2^{-5}} \times \frac{3^{-5}}{3^{-5}} 1 \times 1 \times 5^7$$

$$= 5 \times 5 \times 5 \times 5 \times 5 \times 5 \times 5$$

$$= 78125$$

8. Find x

a. $\left(\frac{2}{5}\right)^{2x+6} \times \left(\frac{2}{5}\right)^3 = \frac{2^{x+2}}{5}$

b. $2^x + 2^x + 2^x = 192$

Sol. a. $\left(\frac{2}{5}\right)^{2x+6} \times \left(\frac{2}{5}\right)^3 = \left(\frac{2}{5}\right)^{x+2}$

$$\left(\frac{2}{5}\right)^{2x+6+3} = \left(\frac{2}{5}\right)^{x+2}$$

$$\left(\frac{2}{5}\right)^{2x+9} = \left(\frac{2}{5}\right)^{x+2}$$

$$2x + 9 = x + 2$$

$$2x - x = 2 - 9$$

$$x = -7$$

b. $2^x + 2^x + 2^x = 192$

$$2^x (1 + 1 + 1) = 192$$

$$2^x = \frac{192}{3}$$

$$2^x = 64$$

$$2^x = (2)^6$$

$$x = 6$$

9. Simplify:

a. $\left(\frac{4}{13}\right)^4 \times \left(\frac{13}{2}\right)^2 \times \left(\frac{7}{4}\right)^3$

b. $\left(\frac{4}{13}\right)^4 \times \left(\frac{13}{2}\right)^2 \times \left(\frac{7}{4}\right)^3$

Sol. a. $\left(\frac{4}{13}\right)^4 \times \left(\frac{13}{2}\right)^2 \times \left(\frac{7}{4}\right)^3$

$$= \frac{4 \times 4 \times 4 \times 4}{13 \times 13 \times 13 \times 13} \times \frac{13 \times 13}{7 \times 7} \times \frac{7 \times 7 \times 7}{4 \times 4 \times 4}$$

$$= \frac{4 \times 7}{13 \times 13} \times \frac{28}{169}$$

b. $\left(\frac{4}{3}\right)^{-2} \times \left(\frac{3}{4}\right)^2 = \left(\frac{3}{4}\right)^2 - \left(\frac{3}{4}\right)^2 = 0$

10. By what number should $\left(-\frac{3}{2}\right)^2$ be divided so that the quotient is $\left(\frac{9}{4}\right)^{-2}$?

Sol. Let the required number = x

According to problem,

$$\left(-\frac{3}{2}\right)^2 \div x = \left(\frac{9}{4}\right)^{-2}$$

$$\text{or } \left(\frac{2}{-3}\right)^3 \div x = \left(\frac{4}{9}\right)^2$$

$$\text{or } \frac{8}{-27} \times x^{-1} = \frac{16}{81} \quad \left[\because \frac{1}{x^a} = x^{-a} \right]$$

$$\text{or } x^{-1} = \frac{16}{81} \times \frac{-27}{8}$$

$$\text{or } x^{-1} = \frac{-2}{3}$$

$$\text{or } \frac{1}{x} = \frac{-2}{3}$$

$$\text{or } \frac{1}{x} = \frac{-2}{3}$$

$$\text{or } x = \frac{-3}{2}$$

$$\text{Hence, required number} = \frac{-3}{2}$$

II. Long answer type questions.

1. Simplify:

$$\text{i. } \left[\left(\frac{2}{3}\right)^{-2}\right]^3 \times \left(\frac{1}{3}\right)^{-4} \times 3^{-1} \times \frac{1}{6}$$

$$\text{ii. } \frac{49 \times z^{-3}}{7^{-3} \times 10 \times z^{-5}} \quad (z \neq 0)$$

$$\text{Sol. i. } \left[\left(\frac{2}{3}\right)^{-2}\right]^3 \times \left(\frac{1}{3}\right)^{-4} \times 3^{-1} \times \frac{1}{6}$$

$$= \left(\frac{-2}{3}\right)^{-6} \times (3^{-1})^{-4} \times 3^{-1} \times \frac{1}{2 \times 3}$$

$$= \left(\frac{-2}{3}\right)^{-6} \times (3)^6 \times 3^4 \times 3^{-1} \times 3^{-1} \times 2^{-1}$$

$$= (2)^{-6-1} \times (3)^{6+4-1-1}$$

$$[\because (-2)^m = (2)^m \text{ if } m \text{ is even}]$$

$$= \frac{(3)^8}{(2)^7} = \frac{3^8}{2^7}$$

$$\text{ii. } \frac{49 \times z^{-3}}{7^{-3} \times 10 \times z^{-5}} \quad (z \neq 0)$$

$$= \frac{7^2 \times 7^3 \times z^{-3}}{10} \times z^5 = \frac{7^5 z^2}{10}$$

2. By what number should $(-8)^{-3}$ be multiplied so that the product may be equal to $(-6)^{-3}$?

[NCERT Exemplar]

Sol. Let the required number be x

$$(-8)^{-3} \times x = (-6)^{-3}$$

$$\left(-\frac{1}{8}\right)^3 \times x = \left(-\frac{1}{6}\right)^3$$

$$x = \left(-\frac{1}{6}\right)^3 \div \left(-\frac{1}{8}\right)^3 = \left(-\frac{1}{6}\right)^3 \times (-8)^3$$

$$= \left(-\frac{1}{6} \times -8\right)^3 \quad (a^m \times b^m = (ab)^m)$$

$$x = \left(\frac{8}{6}\right)^3 = \left(\frac{4}{3}\right)^3 = \frac{64}{27}$$

3. By what number should $\left(\frac{4}{3}\right)^3$ be multiplied, so that the product may be equal to $\left(\frac{-3}{8}\right)^{-3}$?

Sol. Let required number be x .

Now, according to question

$$\left(\frac{1}{2}\right)^3 \times x = \left(\frac{-3}{8}\right)^{-3}$$

Dividing both sides by $\left(\frac{1}{2}\right)^3$, we get

$$\begin{aligned} \Rightarrow \left(\frac{1}{2}\right)^3 \times x \div \left(\frac{1}{2}\right)^3 &= \left(\frac{-3}{8}\right)^{-3} \div \left(\frac{1}{2}\right)^3 \Rightarrow x = \left(\frac{3}{8}\right)^{-3} \div \left(\frac{1}{2}\right)^3 \\ \Rightarrow x &= \left(\frac{8}{-3}\right)^3 \div \left(\frac{1}{2}\right)^3 \quad \left[\because \left(\frac{p}{q}\right)^{-m} = \left(\frac{q}{p}\right)^m \right] \\ \Rightarrow x &= \left(\frac{8}{-3}\right)^3 \div \left(\frac{1}{2}\right)^3 \quad \left[\because (-a)^n = -a^n, \text{ where } n \text{ is odd} \right] \\ \Rightarrow x &= -\frac{8 \times 8 \times 8}{3 \times 3 \times 3} \div \frac{1}{2 \times 2 \times 2} \Rightarrow x = \frac{8 \times 8 \times 8}{3 \times 3 \times 3} \times \frac{2 \times 2 \times 2}{1} \\ \Rightarrow x &= -\left(\frac{16}{3}\right)^3 \end{aligned}$$

4. Mass of Mars is 6.42×10^{29} kg and mass of the Sun is 1.99×10^{30} kg. What is their total mass?

Sol. Given

$$\text{Mass of Mars} = 6.42 \times 10^{29} \text{ kg}$$

$$\text{Mass of the Sun} = 1.99 \times 10^{30} \text{ kg}$$

Converting the standard form into number with the same exponent, we get

$$\text{Mass of Mars} = 6.42 \times 10^{29} \text{ kg}$$

$$\text{Mass of the Sun} = 1.99 \times 10^{30} \text{ kg} = 1.99 \times 10^{29} \times 10 \text{ kg}$$

$$\begin{aligned} \therefore \text{Total mass} &= 6.42 \times 10^{29} \text{ kg} + 1.99 \times 10^{29} \times 10 \text{ kg} \\ &= (6.42 + 1.99 \times 10) \times 10^{29} \text{ kg} \\ &= 26.32 \times 10^{29} \text{ kg} \end{aligned}$$

5. Divide 256 by 10,00,000 and express the result in standard form.

Sol. We have,
$$\frac{256}{10,00,000}$$

$$= 256 \times 10^{-6}$$

$$= 2.56 \times 10^2 \times 10^{-6} = 2.56 \times 10^{-4}$$

I. Higher Order Thinking Skills.

1. By what number should $\left(\frac{-3}{2}\right)^{-3}$ be divided so that the quotient may be $\left(\frac{4}{27}\right)^{-4}$?

Sol. Let the required number be x .

$$\text{Then, } \left(\frac{-3}{2}\right)^{-3} \div x = \left(\frac{4}{27}\right)^{-2}$$

$$\left(\frac{2}{3}\right)^3 \div x = \left(\frac{27}{4}\right)^2$$

$$\frac{8}{-27} \times \frac{1}{x} = \left(\frac{27}{4}\right)^2$$

$$\frac{1}{x} = \frac{-27}{8} \times \frac{(27)^2}{4^2}$$

$$= \frac{-1 \times (27) \times (27)^2}{2 \times (4) \times (4)^2}$$

$$\frac{1}{x} = \frac{-(27)^3}{2(4)^3}$$

$$x = -2 \times \left(\frac{4}{27}\right)^3$$

II. Higher Order Thinking Skills.

1. The size of a red blood cell is 0.000007 m and the size of a plant cell is 0.00001275 m. Compare these two.

Sol. We have,

$$\text{Size of red blood cell} = 0.000007 \text{ m} = 7 \times 10^{-6} \text{ m}$$

$$\text{Size of plant cell} = 0.00001275 = 1.275 \times 10^{-5} \text{ m}$$

$$\therefore \frac{\text{Size of red blood cell}}{\text{Size of plant cell}} = \frac{7 \times 10^{-6}}{1.275 \times 10^{-5}} = \frac{7 \times 10^{-6+5}}{1.275} = \frac{7 \times 10^{-1}}{1.275}$$

$$= \frac{0.7}{1.275} = \frac{0.7}{1.3} = \frac{1}{2}$$

So, a red blood cell is approximately half of a plant cell in size.

2. By what number should $\left(\frac{-3}{2}\right)^{-3}$ be divided so that the quotient may be $\left(\frac{4}{27}\right)^{-2}$?

Sol. Let the required number be x . Then

$$\left(\frac{-3}{2}\right)^{-3} \div x = \left(\frac{4}{27}\right)^{-2}$$

$$\Rightarrow \left(\frac{-3}{2}\right)^{-3} \times \frac{1}{x} = \left(\frac{4}{27}\right)^{-2} \Rightarrow x = \left(\frac{-3}{2}\right)^{-3} \div \left(\frac{4}{27}\right)^{-2}$$

$$\Rightarrow x = \left(\frac{2}{-3}\right)^3 \times \left(\frac{4}{27}\right)^2 = \left[\div \left(\frac{a}{b}\right)^{-n} = \left(\frac{b}{a}\right)^n \right]$$

$$\Rightarrow x = \left(\frac{2}{-3}\right)^3 \times \left(\frac{4}{27}\right)^2 = \left(\frac{2}{-3}\right)^3 \times \left(\frac{2^2}{3^3}\right)^2$$

$$\Rightarrow x = \frac{2^3}{(-3)^3} \times \frac{2^4}{3^6} = \frac{2^{3+4}}{(-3)^3 \times (-3)^6}$$

$$\frac{2^7}{(-3)^9} = \frac{-2^7}{3^9} = \frac{(-2)^7}{3^9}$$

3. What is the value of $\left(x^{\frac{b+c}{c-a}}\right)^{\frac{1}{a-b}} \left(x^{\frac{c+a}{a-b}}\right)^{\frac{1}{b-c}} \left(x^{\frac{a+b}{b-c}}\right)^{\frac{1}{c-a}}$?

Sol.
$$\begin{aligned} & \left(x^{\frac{b+c}{c-a}}\right)^{\frac{1}{a-b}} \left(x^{\frac{c+a}{a-b}}\right)^{\frac{1}{b-c}} \left(x^{\frac{a+b}{b-c}}\right)^{\frac{1}{c-a}} \\ &= \frac{b}{x} + \frac{c}{a} + \frac{a}{b} = \frac{b}{x} \\ &= \frac{b}{x} = \frac{0}{x} \\ &= x^0 = 1 \end{aligned}$$

I. Value Based Questions.

1. (a) Find the value of $\left[\left\{\left(\frac{1}{2}\right)^2\right\}^{-2}\right]^{-1}$

(b) Planet A is at a distance of 9.35×10^6 km from Earth and Planet B is 6.27×10^7 km from Earth. Which Planet is nearer to earth.

Sol. (a)
$$\begin{aligned} \left[\left\{\left(\frac{1}{2}\right)^2\right\}^{-2}\right]^{-1} &= \left[\left(\frac{1}{4}\right)^{-2}\right]^{-1} \\ &= \left[\left(\frac{4}{1}\right)^2\right]^{-1} \quad \left\{\because \frac{1}{x^a} = x^{-a}\right\} \\ &= [16]^{-1} \\ &= \frac{1}{16} \end{aligned}$$

(b) Distance between planet A and earth

$$\begin{aligned} &= 9.35 \times 10^6 \text{ km} \\ &= 0.935 \times 10^7 \text{ km} \end{aligned}$$

Distance between planet B and earth

$$= 6.27 \times 10^7 \text{ km}$$

By changing both distances in the same exponent

We can say planet A is nearer to earth.

2. (a) Find the value of x for which

$$\left(\frac{4}{9}\right)^4 \times \left(\frac{4}{9}\right)^{-7} = \left(\frac{4}{9}\right)^{2x-1}$$

(b) Find the value of $\left(\frac{3}{5}\right)^0 + \left(\frac{125}{129}\right)^0 + \left(\frac{10}{7}\right)^0$

Sol. (a) Since,

$$\left(\frac{4}{9}\right)^4 \times \left(\frac{4}{9}\right)^{-7} = \left(\frac{4}{9}\right)^{2x-1}$$

or
$$\left(\frac{4}{9}\right)^{4+(-7)} = \left(\frac{4}{9}\right)^{2x-1}$$

or $\left(\frac{4}{9}\right)^{-3} = \left(\frac{4}{9}\right)^{2x-1}$

By comparing with power then,

$$-3 = 2x - 1$$

or $2x = -3 + 1$

or $2x = -2$

or $x = -\frac{2}{2} = -1$

Hence, $x = -1$

(b) Since, $\left(\frac{3}{5}\right)^0 + \left(\frac{125}{129}\right)^0 + \left(\frac{10}{7}\right)^0$
 $= 1 + 1 + 1 = 3,$
 $[\because x^0 = 1]$

3. The cells of a bacteria doubles in every 20 min. A scientist begins with a single cell.

(a) Now many cells will be there after

i. 10 h ? ii. 25 h ?

(b) What type of value is depicted by the cells of bacteria ?

Sol. (a) (i) The cell, in bacteria double in every 20 min.

$\therefore$ Number of cells in a bacteria after 20 min. = 2

$\therefore$ Number of cells in a bacteria after 40 min.

$$= 2 \times 2 = 2^2$$

$\therefore$ Number of cells in a bacteria after 1 hr

$$= 2^2 \times 2 = 2^3$$

Number of cells in bacteria after 1 hr 40 min.

$$= 2^4 \times 2 = 2^5$$

Number of cells in bacteria after 2 hrs

$$= 2^5 \times 2 = 2^6$$

$$= (2^3)^2$$

$\therefore$ Number of cells in bacteria after 0 hrs

$$= (2^3)^{25} = 2^{75}$$

(b) The value depicted by the cells of bacteria here is that is double itself after 20 min.

in t hrs by $2^{3 \times t}$.